High Impact Rapid Delivery(HIRD) Supplementary Survey 2007 (District MICS survey)

Urban Information

ΤΟΡΙϹ	INDICATOR	ACCRA	KUMASI	NATIONAL MICS 2006
MORTALITY	Children dead before age 5 (per thousand live births)	46	48	111
	Infants dead before first birthday (per thousand live births)	37	43	71
NUTRITIONAL STATUS	Children < 5 underweight (weight for age) (%)	10	9	18
	Children < 5 stunted (height for age) (%)	9	10	22
	Children < 5 wasted (weight for height) (%)	4	5	5
	Infants with low birthweight (< 2500 g) (%)	9	8	9
SALT IODIZATION	Households using adequately iodized salt (%)	72	40	32
BREASTFEEDING	Exclusive breastfeeding (within first 6 months) (%)	61		54
VITAMIN A	Children receiving vitamin A supplementation within previous six months (%)	76	88	60
IMMUNIZATION	Children fully immunized before 1st birthday (%)	87		64
	Children vaccinated against measles before 1st birthday (%)	92		78
MALARIA MANAGEMENT	Children <5 sleeping under Insectiside treated nets (ITN) (%)	18	19	22
	Households with at least one Insectiside treated net (ITN) (%)	23	27	19
REPRODUCTIVE HEALTH	Married women using contraception (%)	20	11	17
	Women with skilled supervision at delivery (%)	90	92	50
	Women attending at least four antenatal care visits (%)	87	97	n.a.
HIV	Women (15-24 y) with comprehensive knowledge of HIV prevention (%)	37	40	25
EDUCATION	Children of primary school age attending primary school (%)	95	96	75
	Children of secondary school age attending secondary school (%)	68	62	45
	Young women (15-24 y) that are literate (%)	89	83	68
CHILD PROTECTION	Women (20-49 y) married before 18th birthday (%)	13	27	26
	Children with registered births (%)	79	68	51
	Children involved in child labor (%)	15	8	34
WATER AND SANITATION	Access to improved drinking water (%)	98	91	78
	Access to improved sanitation facilities (shared and unshared) (%)	92	96	61
	Percentage of improved facilities that are unshared (%)	17	21	n.a.
National Health Insurance Scheme (NHIS)	Women (15-49 y) registered with NHIS (%)	34	50	n.a.

