


CHILD HEALTH Guyana 2006 Multiple Indicator Cluster Survey (MICS)

| PERCENTAGE OF CHILDREN AGED 0-59 MONTHS WITH DIARRHOEA | ΤΟΡΙΟ | INDICATOR | URBAN COASTAL | RURAL COASTAL | INTERIOR | TOTAL | UNDER-FIVE MORTALITY RATE (died before age 5 years) |
|--|---|---|------------------|------------------|----------|-------|--|
| IN THE TWO WEEKS PRECEDING THE SURVEY AND TREATMENT WITH ORAL REHYDRATION | Child mortality | Under-five mortality rate (died before age 5 years) | 28.0 | 38.0 | 49.0 | 47.0 | 60 1 |
| | | Infant mortality rate (died before first birthday) | 34.0 | 47.0 | 64.0 | 37.0 | 50 - |
| | Nutrition | Underweight (weight for age) | 13.7 | 12.5 | 10.6 | 12.4 | |
| Regions 2,3 Regions 4 Regions 1,7,8,9 Percent 3,4 6,7 9,0 10,5 12,2 Missing Data | | Stunting (height for age) | 11.0 | 11.9 | 21.3 | 13.7 | |
| | | Wasting (reduced weight for height) | 10.8 | 7.9 | 3.3 | 7.6 | |
| | | Low birth weight (< 2,500g) | 17.3 | 17.9 | 23.7 | 18.9 | |
| | Breastfeeding | Timely initiation of breastfeeding (within 1 hour) | 41.5 | 46.4 | 35.1 | 43.1 | CHILD UNDER 5 YEARS OLD HAVING DIARRHOEA IN LAST TWO WEEKS PRECEDING THE SURVEY |
| | | Exclusive breastfeeding rate (under 6 months) | 20.9 | 12.0 | 45.6 | 21.4 | |
| | Immunization | Measles immunization coverage (18-29 months) | 94.8 | 95.4 | 96.4 | 95.4 | |
| | Care of illness | Had diarrhoea in last two weeks | 7.2 | 7.9 | 14.5 | 8.9 | 10 |
| | | Use of oral rehydration therapy (ORT) | 63.5 | 47.7 | 50.8 | 51.7 | |
| | | Had symptoms of pneumonia in last two weeks | 6.5 | 4.1 | 8.8 | 5.6 | |
| | | Care seeking for suspected pneumonia | 60.0 | 59.6 | 74.9 | 64.1 | |
| | | Antibiotic treatment for pneumonia | 40.3 | 14.5 | 6.1 | 19.5 | |
| UNDER-FIVE MORTALITY RATE (died before age 5 years) | Solid fuel use | Households using wood and charcoal for cooking purposes | 2.1 | 9.6 | 47.7 | 10.6 | EXCLUSIVE BREASTFEEDING RATE (under 6 months) |
| | Appropriate water treatment | All drinking water sources (piped, wells, spring, rain, bottled, tanker, cart, surface water) | 55.9 | 51.0 | 29.0 | 49.8 | 30 |
| | | Improved drinking water (piped, wells, spring, rain, bottled) | 56.4 | 51.4 | 35.9 | 51.8 | 20 21 21 21 20 20 20 20 20 20 20 20 20 20 20 20 20 |
| Regions 2,3 | All values relate to children under 5 years of age unless otherwise indicated and all values are in percentages except the mortality figures which are referenced to 1,000 live births. | | | | | | |
| Regions 1,7,8,9 Regions 10 | | | | | | | IMPROVED DRINKING WATER (piped, wells spring, rain, bottled) |
| Number 31 45 | | | | · ·· | | A | |



KEY SURINAME GUYANA TRINIDAD AND TOBAGO BELIZE JAMAICA





47 49 68

Missing Data

